

ISSN 2451-7100

IMAL preprints

<http://www.imal.santafe-conicet.gov.ar/publicaciones/preprints/index.php>

ESTIMATING SUFFICIENT REDUCTIONS OF THE PREDICTORS IN ABUNDANT HIGH-DIMENSIONAL REGRESSIONS

By

R. Dennis Cook, Liliana Forzani and Adam J. Rothman

IMAL PREPRINT # 2012-0006

Publication date: May 8, 2012

Editorial: Instituto de Matemática Aplicada del Litoral
IMAL (CCT CONICET Santa Fe – UNL)
<http://www.imal.santafe-conicet.gov.ar>

Publications Director: Dr. Rubén Spies
E-mail: rspies@santafe-conicet.gov.ar



This article has been published in “*The Annals of Statistics*”, Vol. 40, No. 1, (2012) , pp 353-384.

DOI: [10.1214/11-AOS962](https://doi.org/10.1214/11-AOS962)

See:

<http://projecteuclid.org/DPubS?service=UI&version=1.0&verb=Display&handle=euclid-aos/1333567193>

Please contact the authors for further information.

The Annals of Statistics
2012, Vol. 40, No. 1, 353–384
DOI: 10.1214/11-AOS962
© Institute of Mathematical Statistics, 2012

ESTIMATING SUFFICIENT REDUCTIONS OF THE PREDICTORS IN ABUNDANT HIGH-DIMENSIONAL REGRESSIONS

BY R. DENNIS COOK¹, LILIANA FORZANI AND ADAM J. ROTHMAN²

*University of Minnesota, Instituto de Matemática Aplicada del Litoral
and University of Minnesota*

We study the asymptotic behavior of a class of methods for sufficient dimension reduction in high-dimension regressions, as the sample size and number of predictors grow in various alignments. It is demonstrated that these methods are consistent in a variety of settings, particularly in abundant regressions where most predictors contribute some information on the response, and oracle rates are possible. Simulation results are presented to support the theoretical conclusion.